

Increasing corn for biofuel production reduces biocontrol services in agricultural landscapes

Douglas A. Landis^{a,1}, Mary M. Gardiner^a, Wopke van der Werf^b, and Scott M. Swinton^c

Departments of ^aEntomology and ^cAgricultural, Food, and Resource Economics, Michigan State University, East Lansing MI 48824; and ^bCentre for Crop Systems Analysis, Department of Plant Sciences, Wageningen University, P.O. Box 430, 6700 AK Wageningen, The Netherlands

Edited by May R. Berenbaum, University of Illinois, Urbana, IL, and approved November 5, 2008 (received for review May 22, 2008)

Increased demand for corn grain as an ethanol feedstock is altering U.S. agricultural landscapes and the ecosystem services they provide. From 2006 to 2007, corn acreage increased 19% nationally, resulting in reduced crop diversity in many areas. Biological control of insects is an ecosystem service that is strongly influenced by local landscape structure. Here, we estimate the value of natural biological control of the soybean aphid, a major pest in agricultural landscapes, and the economic impacts of reduced biocontrol caused by increased corn production in 4 U.S. states (Iowa, Michigan, Minnesota, and Wisconsin). For producers who use an integrated pest management strategy including insecticides as needed, natural suppression of soybean aphid in soybean is worth an average of \$33 ha⁻¹. At 2007–2008 prices these services are worth at least \$239 million y⁻¹ in these 4 states. Recent biofuel-driven growth in corn planting results in lower landscape diversity, altering the supply of aphid natural enemies to soybean fields and reducing biocontrol services by 24%. This loss of biocontrol services cost soybean producers in these states an estimated \$58 million y⁻¹ in reduced yield and increased pesticide use. For producers who rely solely on biological control, the value of lost services is much greater. These findings from a single pest in 1 crop suggest that the value of biocontrol services to the U.S. economy may be underestimated. Furthermore, we suggest that development of cellulosic ethanol production processes that use a variety of feedstocks could foster increased diversity in agricultural landscapes and enhance arthropod-mediated ecosystem services.

bioenergy | biological control | ecosystem services

High recent prices of oil and a growing interest in developing alternative liquid fuels has driven a rapid expansion of the corn ethanol industry in the United States. Continuing growth of ethanol production facilities in major corn-producing areas has significantly increased demand for corn grain (1) and is restructuring agricultural landscapes. In 2007, corn plantings in the U.S. totaled 37.9 million ha, a 19% increase over 2006 (ref. 2 and http://usda.mannlib.cornell.edu/usda/nass/CropProd//2000s/2008/CropProd-08-12-2008_revision.pdf). In 2008 corn plantings declined 7% from 2007 levels but still represent the second highest since 1946 (ref. 3 and <http://usda.mannlib.cornell.edu/usda/ers/FDS//2000s/2008/FDS-08-14-2008.pdf>). The Energy Independence and Security Act of 2007 mandates a nearly 5-fold expansion of biofuel production (4), which will likely drive further expansion of corn area. Increases in corn production are already having repercussions, driving up the prices of other major field crops that compete with corn for land. Increased corn acreage for biofuel production has raised concerns regarding the potential for increased food prices, fertilizer and pesticide pollution, soil erosion, biodiversity losses, and greenhouse gas emissions (5–7). Here, we consider a largely unrecognized impact, the effect of a change in landscape structure on arthropod-mediated ecosystem services and its implications for the sustainability of agricultural production systems (8).

It has been estimated that insects provide human society with ecosystem services valued at >\$57 billion yr⁻¹ in the United States (9). Of this, \$4.5 billion yr⁻¹ has been attributed to natural

pest control in agricultural crops. Much of this pest control service is provided by generalist natural enemies that suppress populations of a variety of native and exotic insect pests. Diverse, small-scale agricultural landscapes with a high proportion of noncrop habitats frequently support a greater abundance of natural enemies and lower pest populations than large-scale, monoculture landscapes with little noncrop habitat (10). Expansion of bioenergy crop production on arable lands is likely to change the habitat characteristics that enable these landscapes to support biological control. Increased planting of biofuel crops such as corn, that already dominate large areas in agricultural landscapes, may well reduce biocontrol services.

The soybean aphid (*Aphis glycines* Matsumura) is an invasive insect pest that has become the most significant threat to soybean production in the United States (11). Soybean aphid is consumed by a diversity of natural enemies, including predators and parasitoids (12, 13), that can provide strong top-down regulation of its populations (14), resulting in increased crop yields (15). The natural enemy complex of soybean aphid is currently dominated by generalist predators, in particular ladybird beetles (Coleoptera: Coccinellidae) (16, 17). The main effect of predation on aphid population growth occurs when aphid populations are still small and vulnerable to predation losses (18, 19). Our recent studies show that these predators are responsive to landscape structure and that habitat diversity at the 1.5-km scale surrounding a soybean field is strongly related to the level of soybean aphid suppression. Landscapes with high levels of corn and soybean production had low habitat diversity and significantly reduced biocontrol services in soybean fields (20). Here, we specifically examine how increasing amounts of corn production alters the value of arthropod biological control as an ecosystem service in agricultural landscapes.

Results

Biocontrol Services in Changing Agricultural Landscapes. Annual crop production dominates agricultural landscapes in our study area of Iowa, Michigan, Minnesota, and Wisconsin, and the composition of these landscapes is rapidly changing because of increased demand for corn as a biofuel feedstock (Table 1). In 2007, harvested corn acreage increased by 12%, 20%, 14%, and 17% in Iowa, Michigan, Minnesota, and Wisconsin, respectively, compared with 2006 acreage (2). This increase was primarily at the expense of harvested soybean acreage, which declined by 16%, 13%, 15%, and 20%, respectively. We found that the proportion of corn ($R^2 = 0.66$, $P < 0.0001$) and soybean ($R^2 =$

Author contributions: D.A.L., M.M.G., W.v.d.W., and S.M.S. designed research; M.M.G. and W.v.d.W. performed research; M.M.G. and W.v.d.W. analyzed data; and D.A.L., M.M.G., W.v.d.W., and S.M.S. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. E-mail: landisd@msu.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0804951106/DCSupplemental.

© 2008 by The National Academy of Sciences of the USA




Fig. 3. Value (mean \pm SEM) of biocontrol services against soybean aphid in Iowa, Michigan, Minnesota, and Wisconsin in years of high (2005, $n = 10$ sites) or low (2006, $n = 13$ sites) aphid abundance. Average values represent means across the 2 years. Values shown are based on 2007–2008 projected soybean price of $\$380 \text{ Mg}^{-1}$ for 2 aphid management strategies: biocontrol alone or IPM, under existing landscape conditions in each year and estimated with increased corn production for biofuel based on 2007 changes in corn acres harvested per state.

From this point the model projects continued population growth of aphids until changes in host plant quality cause their decline. By contrast, in the open field with natural enemies present, population growth is greatly suppressed. Using the same initial experimental data, the IPM example yields a markedly different result (Fig. 2B). Here, the observed aphid population on the natural enemy-free plants again rises throughout the initial 14 days, which in the model projection, triggers an insecticide application that reduces the population by 98%. In the presence of natural enemies, the open-field population grows more slowly but eventually exceeds the economic threshold, also triggering an insecticide treatment. The above 4 trajectory projections were replicated for each site-year combination under a landscape change scenario where corn area was increased to 2007 levels for each state. The resulting data were analyzed to determine the overall value of biocontrol services to soybean producers and the impact of landscape change on this ecosystem service under varying aphid infestations and management strategies.

Value of Biocontrol Services. The value of soybean aphid biocontrol services varies with soybean aphid abundance, management strategy, and amount of corn in the landscape. Soybean aphid populations fluctuate from year to year (22), resulting in different values of ecosystem services provided in years of high (2005) versus low (2006) soybean aphid abundance (Fig. 3). In 2005, yield losses were high for growers who relied solely on natural biocontrol. Consequently, the value of the ecosystem service of biological control mitigating these losses was also high, averaging $\$406 \text{ ha}^{-1}$ across the 4 states. An increase of corn in these landscapes to 2007 levels reduced the value of the biocontrol service to an average of $\$199 \text{ ha}^{-1}$. Using the IPM strategy greatly reduced economic losses to soybean aphids; however, IPM users still received an average of $\$35 \text{ ha}^{-1}$ in biocontrol services in 2005. This benefit decreased to an average of $\$29 \text{ ha}^{-1}$ when corn was increased to 2007 levels.

In 2006, soybean aphid abundance was dramatically lower across the study region, resulting in similar biocontrol service values for the IPM and biocontrol approaches (Fig. 3). For growers who rely on biocontrol alone, the presence of natural enemies was worth an average of $\$35 \text{ ha}^{-1}$ across the 4 states. Modeling an increase in corn in these landscapes to 2007 levels reduced the value of the biocontrol service to an average of $\$28 \text{ ha}^{-1}$. In 2006, IPM users received an average of $\$32 \text{ ha}^{-1}$ in

biocontrol services, which decreased to an average of $\$23 \text{ ha}^{-1}$ when corn was increased to 2007 levels.

The value of biocontrol services in soybean depends in part on the number of fields in which aphids exceed the economic threshold and thus trigger costly insecticide applications to protect yield. In 2005, model projections show that 100% of the fields where natural enemies were excluded exceeded the economic threshold. In contrast, where natural enemies were not excluded, 30% of these fields remained below the threshold. In 2006, 62% of fields reached threshold in the absence of natural enemies; however, in the presence of natural enemies no fields required insecticide sprays. Using typical pesticide and application costs of $\$24.5 \text{ ha}^{-1}$ (11), it is clear that the action of natural enemies in preventing the need for insecticide applications is a major factor contributing to the value of the biocontrol service. Additional information on the range of biocontrol service values as influenced by aphid abundance, soybean price, management scenarios, and land-use change are given in *SI Text* and *Tables S1 and S2*.

Aggregate Values. Averaged over the 23 site-year combinations, the value of biocontrol services decreased from $\$196 \text{ ha}^{-1}$ to $\$103 \text{ ha}^{-1}$ as a result of increased corn in the landscape when biological control was the sole pest management strategy (Table S2). Under IPM, the average value of biocontrol services decreased from $\$33 \text{ ha}^{-1}$ to $\$25 \text{ ha}^{-1}$. This change of $\$8 \text{ ha}^{-1}$ amounts to nearly a quarter of average crop chemicals cost of $\$35 \text{ ha}^{-1}$ or 3.4% of total operating costs of $\$235 \text{ ha}^{-1}$ for conventional soybean farmers in the U.S. “heartland” and “northern crescent” states during 2006 (ref. 23 and www.ers.usda.gov/Data/CostsAndReturns). The overall percentage of fields reaching threshold with natural enemies present increased from 30% under the 2005/2006 landscape conditions to an estimated 43% with an increase in corn acreage to the 2007 level.

If all producers used IPM, we estimate the value of natural biocontrol services against soybean aphid in the 4 target states at $\$239 \text{ million y}^{-1}$ or $\$1,620 \text{ farm}^{-1} \text{ y}^{-1}$ under 2007–2008 “biofuel-influenced” soybean prices of $\$380 \text{ Mg}^{-1} = \10.40 Bu^{-1} (Table 2). With corn increased to 2007 levels, the value of biocontrol services in these states declines by an estimated 24%, equivalent to $\$58 \text{ million y}^{-1}$ or $\$390 \text{ farm}^{-1} \text{ y}^{-1}$. Because of increased risk of yield loss, sole reliance on natural biocontrol is a much less common grower strategy. However, if followed by all growers, it would yield aggregate annual biocontrol services against soybean aphid exceeding $\$1.4 \text{ billion y}^{-1}$ in the 4 target states. With corn increased to 2007 levels, the annual value of these services would be reduced by 48% or $\$671 \text{ million y}^{-1}$. Thus, in the world of corn ethanol-driven prices the IPM-based figure of $\$239 \text{ million y}^{-1}$ is a lower bound for the value of these services in these 4 states as some growers realize greater gains by relying entirely on natural biocontrol.

Discussion

Increased corn prices and production provide an immediate profitability benefit to corn growers; however, many of these same growers incur hidden costs to their soybean production because of the attendant landscape change. These impacts extend to neighboring producers and society as a whole through an overall reduction in biocontrol services from these landscapes. From 2006 to 2007, U.S. corn acreage increased dramatically, primarily at the expense of soybean but also from reduced production of minor crops and cultivation of formerly fallow areas such as Conservation Reserve Program acreage. The overall impact of these changes is agricultural landscapes with lower habitat diversity, which has been associated with reduced biological control. A recent metaanalysis found that simple agricultural landscapes had lower abundance of natural enemies (76% of studies) and increased pest pressure (45% of studies)

crop production, and increased communication between scientists, policy makers and the biofuels industry, is critically needed to inform decisions on development of sustainable biofuel production technologies (33).

Materials and Methods

Biocontrol Services Data. Gardiner *et al.* (20) measured the biocontrol service supplied by natural enemies of the soybean aphid, *A. glycines*, in soybean fields across 2 years (2005 and 2006) and 4 states (Iowa, Michigan, Minnesota, and Wisconsin). Within each state, 3–8 sites were studied over the 2 seasons, for a total of 23 site-year combinations. To determine the impact of natural enemies on soybean aphid populations, 2 treatments were compared: an open treatment where natural enemies had full access to aphid-infested soybean plants, and a caged treatment where exclusion cages prevented natural enemies from colonizing plants and consuming aphids. At each location, treatments ($n = 4$ replications) were established when fields reached an average of 10 aphids per plant, and plants in both treatments were manipulated to start with this aphid density at day 0. Aphid counts were made 7 and 14 days after the treatments were established. We have previously established that this time period allows observation of the critical predator impacts while minimizing potential for cage effects (12). Based on these data, we calculated a BSI, which is the relative reduction in aphid density caused by predator access over a period of 14 days:

$$BSI = \frac{\sum_{p=1}^4 (Ac_p - Ao_p)}{Ac_p} \cdot \frac{1}{n}, \quad [1]$$

where Ac is the number of aphids on the caged plant on day 14, Ao is the number of aphids on the open plant on day 14, p is plot, and n is the number of replicates for a given site.

Evaluating Biofuel Landscapes. From 2006 to 2007 harvested corn acreage increased by 12–20% in Iowa, Michigan, Minnesota, and Wisconsin (Table 1). To account for this shift in production, we first determined the corn acreage present within a 1.5-km radius surrounding each of the 23 soybean field sites where BSI was measured in 2005 and 2006. Next, we increased corn acreage for each of these landscapes by the proportional increase recorded for the state in which each individual site was located. Finally, we used the equation from the linear regression of corn and BSI ($y = -1.301x + 1.109$) to determine new predicted BSI values for each site based on the increased corn acreage. These BSI values allow for the calculation of adjusted aphid abundance values for the open, i.e., predator-exposed treatment at 14 days of the experiment. We then compare the service provided by natural enemies under 2 landscape regimes, the 2005–2006 landscape (based on land cover data collected in 2005 and 2006) and a biofuels-influenced landscape (incorporating the 2007 corn increase).

Predicting Aphid Population Growth. To predict the value of biocontrol services obtained under both landscape regimes, it was necessary to project aphid population growth beyond the timeframe of the 14-day experiment. To accomplish this we used an aphid population growth model (21) that is based on the growth of natural enemy-free (i.e., caged) soybean aphid populations. We used a linear interpolation to estimate aphid populations during the experimental period 0–14 days then projected the population for the remainder of the season by using Eq. 2:

$$N_t = N_0 e^{rt \left(1 - \frac{1}{2}at\right)}, \quad [2]$$

where N is the aphid population size, t is thermal time since sowing (degree days base = 10 °C), r is the intrinsic rate of increase of soybean aphid at thermal time 0, and a expresses the effect of plant age (thermal time) on the rate of population increase of soybean aphid on soybean plants (34). This model showed good to excellent predictions of soybean aphid population dynamics in the absence of predators with $R^2_{\text{prediction}}$ from independent data ranging from 0.86 to 0.99 (21). The population projection after the 2-week period of the field measurements does not account for further predation effects and therefore provides a lower bound for predator impact.

Soybean Aphid Pest Management Scenarios. We used 2 scenarios to describe producer responses to soybean aphid infestations. The first was IPM, where

producers scout soybean aphid populations every 7 days and make insecticide applications 4 days after a population reaches the economic threshold. Our second strategy was a biocontrol-alone strategy, where producers rely completely on natural biocontrol. IPM strategies represent profit-maximizing behavior for those soybean growers that use pesticides. The economic threshold for pest control is the level at which a pest population will cause crop yield losses that exceed the cost of control, in this case, insecticide treatment. The established economic threshold is 250 aphids per soybean plant (11). Because many producers rely on custom applicators for field spraying services, the 4-day delay represents a realistic average response time. We assume insecticide applications induce 98% mortality and prevent aphid population growth for 7 days.

Estimating Soybean Damage. To estimate the damage caused by soybean aphid in the presence and absence of natural enemies, cumulative aphid days (CAD) were calculated by integrating the area under each curve (aphid population with and without predation) by application of the trapezoidal rule over daily time intervals. To translate CAD into yield loss, we applied a yield loss coefficient of 0.688% per 1,000 aphid days per plant, derived from yield loss studies from 19 location-year combinations in 6 Midwestern states, including all 4 states included in our study (linear regression of yield loss on CAD, $R^2 = 0.665$) (11). To avoid crop loss >100%, the proportional crop loss was truncated at a value of 1.

Calculation of Value of Biocontrol Services. To calculate the monetary value of yield loss caused by soybean aphid we used the 10-year median yields in $\text{Mg}\cdot\text{ha}^{-1}$ from 1997 to 2006 for Iowa (3.16), Michigan (2.59), Minnesota (2.79), and Wisconsin (2.96) (35). Pesticide application costs of $\$24.5\text{ ha}^{-1}$ represent a midrange control cost representative of hired ground application of a moderately-priced soybean aphid insecticide (11). For each year and field we estimated the value of income loss ha^{-1} caused by soybean aphid with or without natural enemies, including yield loss, insecticide, and scouting costs for IPM strategies ($\$5\text{ ha}^{-1}$) (36). We calculated the mean for each set of $n = 23$ site-years per scenario. By subtracting the mean loss with natural enemies present from the loss in their absence, we obtained the value of biocontrol services (VBS) for that scenario. The loss ha^{-1} was multiplied by the area of soybean harvested in each state to provide a statewide estimate of VBS. The difference in VBS ha^{-1} between 2005–2006 and 2007 landscapes was used to calculate the reduced VBS in a biofuel landscape that was scaled to the state level.

Monetary Valuation Method. We adopted the production function approach to monetary valuation (37), which allows the value of biocontrol services to be inferred from their effects on the production of a marketed product, such as soybean. Critics of prior attempts to place monetary values on ecosystem services have highlighted the importance of focusing on marginal changes, consideration of most likely alternatives, and recognition of market price feedback effects (38, 39). For followers of the IPM strategy, the value of biocontrol services comes from both avoiding insecticide application costs and averting yield loss. For producers relying solely on biocontrol, the value of biocontrol services comes entirely from reduced yield loss. Although the relative size of these grower groups is not well known, U.S. Department of Agriculture surveys in 2005–2006 in the 4 states studied found the percentage of soybean acreage with insecticide use ranges from 4% (Wisconsin) to 56% (Minnesota), with a 2-year median of 23% (ref. 40 and <http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1560>). In contrast, only 0.2% of U.S. soybean land is certified for organic production (ref. 41 and www.ers.usda.gov/Data/Organic) and hence committed to a strategy like sole reliance on biocontrol. Given the current situation, we judge the IPM scenario to be the most representative strategy for purposes of valuation of biocontrol as an ecosystem service.

The importance of market-price feedback effects is evident from the recent fluctuations in agricultural commodity prices. Soybean prices more than doubled from their recent historic levels of 1997–2006 to early 2008, when soybean futures market prices reached levels considerably higher than the 2007–2008 U.S. Department of Agriculture price projection of $\$380\text{ Mg}^{-1} = \10.40 Bu^{-1} (ref. 42 and <http://usda.mannlib.cornell.edu/usda/ers/OCS/2000s/2008/OCS-02-11-2008.pdf>). In response, U.S. farmers reduced corn planted area by 7% in 2008 (still the second-highest level since 1946) in favor of more soybean (2). To balance these positive and negative price feedback effects, we contrasted 2 market values for soybean, the current, biofuel-driven price (the U.S. Department of Agriculture 2007–2008 projection) and the recent past 10-year median U.S. soybean prices in 1997–2006 (ref. 42 and <http://usda.mannlib.cornell.edu/usda/ers/OCS/2000s/2008/OCS-02-11-2008.pdf>). Changes in the value of biocontrol services were also contrasted to chemical

and total operating cost for soybean producers in our region (ref. 23 and www.ers.usda.gov/Data/CostsAndReturns).

ACKNOWLEDGMENTS. We thank Steve Scott for programming assistance and Risk Avoidance and Mitigation Program collaborators for supplying the original data and G. Philip Robertson, Douglas Schemske, James R. Miller, Bruce E. Dale, Holger Meinke, Rufus Isaacs, and 3 anonymous reviewers for useful comments that improved the manuscript. This work was supported by the U.S.

Department of Agriculture Cooperative State Research, Education, and Extension Service Risk Avoidance and Mitigation Program Grant 2004-51101-02210), Department of Energy Great Lakes Bioenergy Research Center Grant DE-FC02-07ER64494), the National Science Foundation Long-Term Ecological Research Program at the W.K. Kellogg Biological Station (through Grant 0423627), and the Michigan Agricultural Experiment Station. W.v.d.W. was supported by the C.T. de Wit Graduate School for Production Ecology and Resource Conservation.

- Westcott PC (2007) *Ethanol Expansion in the United States: How Will the Agricultural Sector Adjust?* (Department of Agriculture Economic Research Service, Washington DC).
- US Department of Agriculture (2008) *Crop Production 2007 Summary* CrPr2-2(8-08). (National Agricultural Statistics Service, Washington DC).
- Baker A, Allen E, Lutman H (2008). *Feed Outlook FDS-08g*. (Department of Agriculture Economic Research Service, Washington, DC).
- Rahall N, et al. (2007) *HR 6: Energy Independence and Security Act of 2007* (Library of Congress, Washington, DC).
- Nash S (2007) Decrypting biofuel scenarios. *BioScience* 57:472–477.
- Searchinger T, et al. (2008) Use of U.S. croplands for biofuels increases greenhouse gases through emissions from land-use change. *Science* 319:1238–1240.
- Donner SD, Kucharik CJ (2008) Corn-based ethanol production compromises goal of reducing nitrogen export by the Mississippi River. *Proc Natl Acad Sci USA* 105:4513–4518.
- Swinton SM, Lupi F, Robertson GP, Landis DA (2006) Ecosystem services from agriculture: Looking beyond the usual suspects. *Am J Agric Econ* 88:1160–1166.
- Losey J, Vaughan M (2006) The economic value of ecological services provided by insects. *BioScience* 56:311–323.
- Bianchi FJJA, Booi CJH, Tscharrnke T (2006) Sustainable pest regulation in agricultural landscapes: A review on landscape composition, biodiversity, and natural pest control. *Proc R Soc London Ser B* 273:1715–1727.
- Ragsdale DW, et al. (2007) Economic threshold for soybean aphid (Hemiptera: Aphididae). *J Econ Entomol* 100:1258–1267.
- Fox TB, Landis DA, Cardoso FF, DiFonzo CD (2004) Predators suppress *Aphis glycines* Matsumura population growth in soybean. *Environ Entomol* 33:608–618.
- Rutledge CE, O'Neil RJ, Fox TB, Landis DA (2004) Soybean aphid predators and their use in IPM. *Ann Entomol Soc Am* 97:240–248.
- Costamagna AC, Landis DA (2006) Predators exert top-down control of soybean aphid across a gradient of agricultural management systems. *Ecol Appl* 16:1619–1628.
- Costamagna AC, Landis DA, DiFonzo CD (2007) Suppression of soybean aphid by generalist predators results in a trophic cascade in soybeans. *Ecol Appl* 17:441–451.
- Costamagna AC, Landis DA (2007) Quantifying predation on soybean aphid through direct field observations. *Biol Control* 42:16–24.
- Costamagna AC, Landis DA, Brewer MJ (2008) The role of natural enemy guilds in *Aphis glycines* suppression. *Biol Control* 45:368–379.
- Landis DA, van der Werf W (1997) Early-season aphid predation impacts establishment and spread of sugar beet yellows virus in the Netherlands. *Entomophaga* 42:499–516.
- Ives AR, Settle WH (1997) Metapopulation dynamics and pest control in agricultural systems. *Am Nat* 149:220–246.
- Gardiner MM, et al. (2008) Landscape diversity enhances the biological control of an introduced crop pest in the north-central U.S. *Ecol Appl*, in press.
- Costamagna AC, van der Werf W, Bianchi FJJA, Landis DA (2007) An exponential growth model with decreasing *r* captures bottom-up effects on the population growth *Aphis glycines* Matsumura (Hemiptera: Aphididae). *Agr Forest Entomol* 9:297–305.
- Donaldson JR, Myers SW, Gratton C (2007) Density-dependent responses of soybean aphid (*Aphis glycines* Matsumura) populations to generalist predators in mid- to late-season soybean fields. *Biol Control* 43:111–118.
- US Department of Agriculture (2008) *Conventional Soybean Production Costs and Returns per Planted Acre, by Region, Excluding Government Payments, 2006* (Economic Research Service, Washington DC).
- Maredia KM, Gage SH, Landis DA, Wirth TM (1992) Ecological observations on predator Coccinellidae (Coleoptera) in southwestern Michigan. *Great Lakes Entomol* 25:265–270.
- Colunga-Garcia M, Gage SH, Landis DA (1997) Response of an assemblage of Coccinellidae (Coleoptera) to a diverse agricultural landscape. *Environ Entomol* 26:797–804.
- Barducci TB (1972) in *The Careless Technology: Ecology and International Development*, eds Farvar MT, Milton JP (Natural History Press, Garden City, NY), pp 423–438.
- Schmer MR, Vogel KP, Mitchell RB, Perrin RK (2008) Net energy of cellulosic ethanol from switchgrass. *Proc Natl Acad Sci USA* 105:464–469.
- Tilman D, Hill J, Lehman C (2006) Carbon-negative biofuels from low-input high-diversity grassland biomass. *Science* 314:1598–1600.
- Perlack RD, et al. (2005) *Biomass as a Feedstock for a Bioenergy and Bioproducts Industry: The Technical Feasibility of a Billion-Ton Annual Supply* (Oak Ridge National Laboratory, Oak Ridge, TN).
- Murray LD, Best LB, Jacobsen TJ, Braster M (2003) Potential effects on grassland birds of converting marginal cropland to switchgrass biomass production. *Biomass Bioenergy* 25:167–175.
- McLaughlin SB, Kszos LA (2005) Development of switchgrass (*Panicum virgatum*) as a bioenergy feedstock in the United States. *Biomass Bioenergy* 28:515–535.
- Isaacs R, Tuell J, Fiedler A, Gardiner M, Landis D (2009) Maximizing arthropod-mediated ecosystem services in agricultural landscapes: The role of native plants. *Front Ecol Environ*, 10.1890/080035.
- Robertson GP, et al. (2008) Sustainable biofuels redux. *Science* 322:49–50.
- Williams IS, van der Werf W, Dewar AM, Dixon AFG (1997) Factors affecting the relative abundance of two coexisting aphid species on sugar beet. *Agric Forest Entomol* 1:119–125.
- U.S. Department of Agriculture (1999, 2002, 2005, 2008) *Crop Production Summary Reports* (National Agricultural Statistics Service, Washington DC).
- Song F, Swinton SM, DiFonzo C, O'Neal M, Ragsdale DW (2006) *Staff Paper 2006-24, Department of Agricultural Economics* (Michigan State University, East Lansing).
- Freeman AM (2003) *The Measurement of Environmental and Resource Values: Theory and Methods* (Resources for the Future, Washington DC), 2nd Ed, pp 259–267.
- Bockstael NE, Freeman AM, Kopp RJ, Portney PR, Smith VK (2000) On measuring economic values for nature. *Environ Sci Tech* 34:1384–1389.
- Pearce D (1998) Auditing the earth. *Environment* 40:23–28.
- US Department of Agriculture (2006/2007) *Agricultural Chemical Usage Field Crops Summary* (National Agricultural Statistics Service, Washington DC).
- US Department of Agriculture (2008) *Data Sets: Organic Production* (Department of Agriculture Economic Research Service, Washington DC).
- Ash M, Dohlmán E (2008) *Oil Crops Outlook* (Department of Agriculture Economic Research Service, Washington, DC).